

THE H-N FILTRATION OF BUNDLES AS FROBENIUS PULL-BACK

MINGSHUO ZHOU

ABSTRACT. Let X be a smooth projective curve of genus $g \geq 2$ over an algebraic closed field k of characteristic $p > 0$. Let $F : X \rightarrow X_1$ be the relative Frobenius morphism, and E be a semistable torsion free sheaf on X . For a semistable vector bundle E , one may guess that the length of Harder-Narasimhan filtration is not more than p . In this paper, I give a negative answer to this by giving an explicit example.

1. INTRODUCTION

Let X be a smooth projective curve of genus $g \geq 2$ defined over an algebraic closed field k of characteristic $p > 0$. The absolute Frobenius morphism $F_X : X \rightarrow X$ is induced by $\mathcal{O}_X \rightarrow \mathcal{O}_X, f \mapsto f^p$. Let $F : X \rightarrow X_1 := X \times_k k$ denote the relative Frobenius morphism over k . One of the themes is to study its action on the geometric objects on X .

Recall that a vector bundle E on a smooth projective curve is called semistable (resp. stable) if $\mu(E') \leq \mu(E)$ (resp. $\mu(E') < \mu(E)$) for any nontrivial proper subbundle $E' \subset E$, where $\mu(E)$ is the slope of E . Semistable bundles are basic constituents of vector bundle in the sense that any bundle E admits a unique filtration

$$HN_{\bullet}(E) : 0 = HN_0(E) \subset HN_1(E) \subset \cdots \subset HN_{\ell}(E) = E'$$

which is the so called Harder-Narasimhan filtration, such that

- (1) $gr_i^{HN}(E) := HN_i(E)/HN_{i-1}(E)$ ($1 \leq i \leq \ell$) are semistable;
- (2) $\mu(gr_1^{HN}(E)) > \mu(gr_2^{HN}(E)) > \cdots > \mu(gr_{\ell}^{HN}(E))$. The integer ℓ is called the length of the Harder-Narasimhan filtration of bundle E . It measures how far is a vector bundle from being semistable in some sense, and it's clear E is semistable if and only if $\ell = 1$. It is known that F_* preserves the stability of vector bundles ([6]), but F^* does not preserve the semistability of vector bundle ([3] for example).

Date: April 20, 2012.

The author is supported by gucas.

Given a semistable vector bundle E on X , then F^*E may not be semistable, it's natural to consider the length of the Harder-Narasimhan filtration of F^*E . If $E = F_*W$ where W is a semistable bundle on X , the length of Harder-Narasimhan filtration of F^*E is p by Sun's Theorem (cf. [6, Theorem 2.2]) is p . In this case, the instability of F^*E is $(p-1)(2g-2)$ which reaches the upper bound of F^*E . Actually, in [5], the authors prove the instability $I(F^*E) \leq (p-1)(2g-2)$ for a semistable bundle E on X . In consideration of the above statement, one may guess that $\forall$ semistable bundle E on X , the length of the Harder-Narasimhan filtration of F^*E is not more than p .

In this note, we give a negative answer to the above guess by constructing an example. Briefly, give a polarized smooth projective surface (Y, H) with $\mu(\Omega_Y^1) > 0$ is semistable, for a suitable semistable vector bundle W on Y (take W to be line bundle for example), F_*W is semistable and F^*F_*W has the Harder-Narasimhan filtration whose length is bigger than p . Restricting F^*F_*W and the Harder-Narasimhan filtration to a generic hyperplane X of high degree, by some analysis, one can check it's actually the Harder-Narasimhan filtration of $F^*(E|_X)$, where $E = F_*W$ and the stability of $E|_X$ would be proved in the text.

2. CONSTRUCTION OF THE EXAMPLE

Let Y be a smooth projective variety of dimension n over an algebraic closed field k with $\text{char}(k) = p > 0$. Fix an ample divisor H on Y , for a torsion free sheaf E , the slope of E is defined as

$$\mu(E) = \frac{c_1(E) \cdot H^{n-1}}{\text{rk}(E)}$$

where $\text{rk}(E)$ denotes the rank of E . Then

Definition 2.1. A torsion free sheaf E on Y is called semistable (resp. stable) if for any subsheaf $E' \subset E$ with E/E' torsion free, we have

$$\mu(E') \leq (\text{resp. } <) \mu(E).$$

Theorem 2.2. (Harder-Narasimhan filtration) For any torsion free sheaf E , there is a unique filtration

$$HN_\bullet(E) : 0 = HN_0(E) \subset HN_1(E) \subset \cdots \subset HN_\ell(E) = E'$$

which is the so called Harder-Narasimhan filtration, such that

- (1) $gr_i^{HN}(E) := HN_i(E)/HN_{i-1}(E)$ ($1 \leq i \leq \ell$) are semistable;
- (2) $\mu(gr_1^{HN}(E)) > \mu(gr_2^{HN}(E)) > \cdots > \mu(gr_\ell^{HN}(E))$.

Remark 2.3. In [4, Theorem 1.3.4], the proof of existence of the filtration is given in terms of Gieseker stability. In particular, $gr_i^{HN}(E)$ are Gieseker semistable, thus they are μ semistable torsion free sheaves. We call the integer ℓ is the length of the Harder-Narasimhan filtration.

Definition 2.4. Let E be a semistable sheaf. A Jordan-Hölder filtration of E is a filtration

$$0 = E_0 \subset E_1 \subset \cdots \subset E_\ell = E$$

such that the factors $gr_i(E) = E_i/E_{i-1}$ are stable .

Remark 2.5. Jordan-Hölder filtration always exists. As in Remark 2.3, we can get the factors $gr_i(E)$ are torsion free.

Let $F : Y \rightarrow Y_1 := Y \times_k k$ denote the relative Frobenius morphism over k .

Lemma 2.6. *Let X be a closed subvariety of Y , then the induced morphism $F|_C : C \rightarrow C_1$ is the relative Frobenius morphism.*

Proof. For the case of absolute Frobenius morphism, it's trivial. In the relative case, it's just a translation of the absolute case to the relative case. $\square$

Now, let's restrict Y to be a smooth projective variety of dimension n with Ω_Y^1 is semistable and $\mu(\Omega_Y^1) > 0$ (For example, Y can be chosen to be $C \times C$ with C is a smooth projective curve with genus $g \geq 2$). Let W be a torsion free sheaf on Y , define $V_0 := V = F^*(F_*W)$, $V_1 = \ker(F^*(F_*W)) \rightarrow W$,

$$V_{\ell+1} := \ker((V_\ell) \xrightarrow{\nabla} V \otimes_{\mathcal{O}_Y} \Omega_Y^1 \rightarrow (V/V_\ell) \otimes_{\mathcal{O}_Y} \Omega_Y^1)$$

where $\nabla : V \rightarrow V \otimes_{\mathcal{O}_Y} \Omega_Y^1$ is the canonical connection. Actually, for the above filtration we have

Theorem 2.7. ([6, Theorem 3.7]) *The filtration defined above is*

$$0 = V_{n(p-1)+1} \subset V_{n(p-1)} \subset \cdots \subset V_1 \subset V_0 = V = F^*(F_*W)$$

which has the following properties

- (1) $\nabla(V_{\ell+1}) \subset V_\ell \otimes_{\mathcal{O}_Y} \Omega_Y^1$ for $\ell \geq 1$, and $V_0/V_\ell \cong W$.
- (2) $V_\ell/V_{\ell+1} \xrightarrow{\nabla} (V_{\ell-1}/V_\ell) \otimes_{\mathcal{O}_Y} \Omega_Y^1$ are injective for $1 \leq \ell \leq n(p-1)$, which induced isomorphisms

$$\nabla^\ell : V_\ell/V_{\ell+1} \cong W \otimes_{\mathcal{O}_Y} T^\ell(\Omega_Y^1), \quad 0 \leq \ell \leq n(p-1)$$

The vector bundle $T^\ell(\Omega_Y^1)$ is suited in the exact sequence

$$\begin{aligned} 0 \rightarrow \text{Sym}^{\ell-\ell(p) \cdot p}(\Omega_Y^1) \otimes F^* \Omega_Y^{\ell(p)} &\xrightarrow{\phi} \text{Sym}^{\ell-(\ell(p)-1) \cdot p}(\Omega_Y^1) \otimes F'^* \Omega_Y^{\ell(p)-1} \\ &\rightarrow \cdots \rightarrow \text{Sym}^{\ell-q \cdot p}(\Omega_Y^1) \otimes F^* \Omega_Y^q \xrightarrow{\phi} \text{Sym}^{\ell-(q-1) \cdot p}(\Omega_Y^1) \otimes F'^* \Omega_Y^{q-1} \\ &\rightarrow \cdots \rightarrow \text{Sym}^{\ell-p}(\Omega_Y^1) \otimes F^* \Omega_Y^1 \xrightarrow{\phi} \text{Sym}^\ell(\Omega_Y^1) \rightarrow T^\ell(\Omega_Y^1) \rightarrow 0 \end{aligned}$$

where $\ell(p) \geq 0$ is the integer such that $\ell - \ell(p) \cdot p < p$.

Lemma 2.8. *If Ω_Y^1 is strongly semistable with $\mu(\Omega_Y^1) > 0$ and W is strongly semistable, then the filtration defined in Theorem 2.7*

$$0 = V_{n(p-1)+1} \subset V_{n(p-1)} \subset \cdots \subset V_1 \subset V_0 = V = F^*(F_*W)$$

is the Harder-Narasimhan filtration of $F^*(F_*W)$.

Proof. From the Lemma 4.5 of [7], one can get

$$\mu(V_\ell/V_{\ell+1}) = \frac{l}{n} K_Y \cdot H^{n-1} + \mu(W).$$

So the rest is enough to show that $W \otimes_{\mathcal{O}_Y} T^\ell(\Omega_Y^1)$ is semistable. By W is strongly semistable, it's enough to show $T^\ell(\Omega_Y^1)$ is strongly semistable. In the long exact sequence of Theorem 2.7, every terms have slope $\ell \cdot \mu(\Omega_Y^1)$ by direct computations and strongly semistable by easily checking. Now, the lemma comes from the following fact: For a short exact sequence of torsion free sheaves with the same slope:

$$0 \rightarrow E_1 \rightarrow E_2 \rightarrow E_3 \rightarrow 0,$$

then one of them is strongly semistable if the other two is strongly semistable. $\square$

Remark 2.9. Under the above condition, one can deduce F_*W is semistable, denote it by E . Actually, it is a direct corollary of the following Theorem.

Theorem 2.10. ([7, Theorem 4.2]) *When $K_Y \cdot H^{n-1} \geq 0$, we have, for any $E \subset F_*W$,*

$$\mu(F_*W) - \mu(E) \geq -\frac{I(W, X)}{P}.$$

In particular, if $W \otimes T^\ell(\Omega_Y^1)$, $0 \leq \ell \leq n(p-1)$, are semistable, then F_*W is semistable. Moreover, if $K_Y \cdot H^{n-1} > 0$, the stability of the bundles $W \otimes T^\ell(\Omega_Y^1)$, $0 \leq \ell \leq n(p-1)$, implies the stability of F_*W .

Lemma 2.11. ([2, Corollary 5.4]) *Let E be a torsion-free sheaf of rank $r > 1$ on Y . Assume that E is μ semistable with respect to $(D_1, \dots, D_{n-1})$ and let $0 = E_0 \subset E_1 \subset \cdots \subset E_m = E$ be the corresponding Jordan-Hölder filtration of E , set $F_i = E_i/E_{i-1}$, $r_i = \text{rk} F_i$.*

Let $D \in |kD_1|$ be a normal divisor such that all the sheaves $F_i|_D$ have no torsion. If

$$k > \lfloor \frac{r-1}{r} \Delta(E) D_2 \cdots D_{n-1} + \frac{1}{dr(r-1)} + \frac{(r-1)\beta_r}{dr} \rfloor$$

then $E|_D$ is μ semistable with $(D_2|_D, \dots, D_{n-1}|_D)$.

Remark 2.12. For the notation, one can see [2] for detail. Note that if E is torsion free then the restriction $E|_D$ is also torsion free for a general divisor D in a base point-free system (see [4, Corollary 1.1.14] for a precise statement). Take $k \gg 0$, using the Bertini's Theorem (cf. [1, Theorem 8.8]), we can choose $D \in |kD_1|$ to be smooth projective variety and $E|_D$ is a semistable torsion free sheaf with respect to $(D_2|_D, \dots, D_{n-1}|_D)$. Repeating the above process, we can get a smooth projective curve X which is a subvariety of Y , such that $E|_C$ is a semistable vector bundle. From our choice, the curve C has large genus.

Now, we can construct the example using the above preliminaries.

Construction 2.13. Let Y to be a smooth projective variety of dimension n with Ω_Y^1 is strongly semistable and $\mu(\Omega_Y^1) > 0$, for example we can take $Y = C \times \cdots \times C$ with C is a smooth curve of genus $G \geq 2$. Let W to be a strongly semistable bundle on Y , for example we can take W to be the copies of line bundle. Consider the following diagram:

$$\begin{array}{ccc} Y & \xrightarrow{F} & Y \\ \uparrow & & \uparrow \\ X & \xrightarrow{F} & X \end{array}$$

where X is smooth projective curve which is chosen as in Remark 2.13, and the commutative diagram comes from Lemma 2.6. Denote F_*W by E , then E is semistable from Remark 2.9 and $E|_X$ is semistable from Remark 2.12. Denote the Harder-Narasimhan filtration of F^*E

$$0 = V_{n(p-1)+1} \subset V_{n(p-1)} \subset \cdots \subset V_1 \subset V_0 = V = F^*(F_*W)$$

by **HN**, we have the length of **HN** is $n(p-1)+1$. Consider the restriction of the filtration HN to X which denoted by $\mathbf{HN}|_X$, $(V_i/V_{i+1})|_X = (V_i|_X)/(V_{i+1}|_X)$ is semistable by Remark 2.12 and $\mu((V_i|_X)/(V_{i+1}|_X))$ is strictly increasing. Meanwhile, $(F^*E)|_X = F^*(E|_X)$ by the commutative diagram and $\mathbf{HN}|_X$ is the Harder-Narasimhan filtration of $F^*(E|_X)$. Above all, we construct a semistable bundle $E|_X$ over a smooth curve X with genus $g > 2$, such that the length of the Harder-Narasimhan filtration is $n(p-1)+1$. So there is no bound for the

length of the Harder-Narasimhan filtration of a bundle as Frobenius pull-back of a semistable bundle.

REFERENCES

- [1] R. Hartshorne: *Algebraic geometry*, Berlin-Heidelberg-New York. Springer, 1977.
- [2] A.langer: *Semistable sheaves in positive characteristic*, Ann.Math. **159** (2004),251-276.
- [3] D.Gieseker: *Stable vector bundles and the Frobenius morphism*, Ann.Sci.École Norm.Sup. (4),**6**(1973),95-101.
- [4] D.Huybrechts and M.Lehn, *The Geometry of Moduli Spaces of Sheaves*, Aspects of Mathematics **31**, Friedr. Vieweg & Sohn, Braunschweig (1977).
- [5] V.Mehta and C.pauly: *Semistability of Frobenius direct images over curves*, Bull. SOC. Math. France, **135** (2007), 105-117.
- [6] X. Sun: *Direct images of bundles under Frobenius morphism*, Invent. math. **173** (2008), 427-447.
- [7] X. sun: *Frobenius morphism and semistable bundles*, Advanced Studies in Pure Mathematics 60 (2010), Algebraic Geometry in East Asia- Seoul (2008), !61-182.

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCE, BEIJING, P. R. OF CHINA

E-mail address: zhoumingshuo@amss.ac.cn